

APPENDICES

Appendix I. Complex Numbers (page A-10)

1. $z = -5 + 2i$, $\operatorname{Re}(z) = -5$, $\operatorname{Im}(z) = 2$




Fig. .1

2. $z = 4 - i$, $\operatorname{Re}(z) = 4$, $\operatorname{Im}(z) = -1$

3. $z = -\pi i$, $\operatorname{Re}(z) = 0$, $\operatorname{Im}(z) = -\pi$

4. $z = -6$, $\operatorname{Re}(z) = -6$, $\operatorname{Im}(z) = 0$

5. $z = -1 + i$, $|z| = \sqrt{2}$, $\operatorname{Arg}(z) = 3\pi/4$
 $z = \sqrt{2}(\cos(3\pi/4) + i \sin(3\pi/4))$

6. $z = -2$, $|z| = 2$, $\operatorname{Arg}(z) = \pi$
 $z = 2(\cos \pi + i \sin \pi)$

7. $z = 3i$, $|z| = 3$, $\operatorname{Arg}(z) = \pi/2$
 $z = 3(\cos(\pi/2) + i \sin(\pi/2))$

8. $z = -5i$, $|z| = 5$, $\operatorname{Arg}(z) = -\pi/2$
 $z = 5(\cos(-\pi/2) + i \sin(-\pi/2))$

9. $z = 1 + 2i$, $|z| = \sqrt{5}$, $\theta = \operatorname{Arg}(z) = \tan^{-1}2$
 $z = \sqrt{5}(\cos \theta + i \sin \theta)$

10. $z = -2 + i$, $|z| = \sqrt{5}$, $\theta = \operatorname{Arg}(z) = \pi - \tan^{-1}(1/2)$
 $z = \sqrt{5}(\cos \theta + i \sin \theta)$

11. $z = -3 - 4i$, $|z| = 5$, $\theta = \operatorname{Arg}(z) = -\pi + \tan^{-1}(4/3)$
 $z = 5(\cos \theta + i \sin \theta)$

12. $z = 3 - 4i$, $|z| = 5$, $\theta = \operatorname{Arg}(z) = -\tan^{-1}(4/3)$
 $z = 5(\cos \theta + i \sin \theta)$

13. $z = \sqrt{3} - i$, $|z| = 2$, $\operatorname{Arg}(z) = -\pi/6$
 $z = 2(\cos(-\pi/6) + i \sin(-\pi/6))$

14. $z = -\sqrt{3} - 3i$, $|z| = 2\sqrt{3}$, $\operatorname{Arg}(z) = -2\pi/3$
 $z = 2\sqrt{3}(\cos(-2\pi/3) + i \sin(-2\pi/3))$

15. $z = 3 \cos \frac{4\pi}{5} + 3i \sin \frac{4\pi}{5}$
 $|z| = 3$, $\operatorname{Arg}(z) = \frac{4\pi}{5}$

16. If $\operatorname{Arg}(z) = \frac{3\pi}{4}$ and $\operatorname{Arg}(w) = \frac{\pi}{2}$, then
 $\arg(zw) = \frac{3\pi}{4} + \frac{\pi}{2} = \frac{5\pi}{4}$, so
 $\operatorname{Arg}(zw) = \frac{5\pi}{4} - 2\pi = \frac{-3\pi}{4}$.

17. If $\operatorname{Arg}(z) = -\frac{5\pi}{6}$ and $\operatorname{Arg}(w) = \frac{\pi}{4}$, then
 $\arg(z/w) = -\frac{5\pi}{6} - \frac{\pi}{4} = -\frac{13\pi}{12}$, so
 $\operatorname{Arg}(z/w) = -\frac{13\pi}{12} + 2\pi = \frac{11\pi}{12}$.

18. $|z| = 2$, $\arg(z) = \pi \Rightarrow z = 2(\cos \pi + i \sin \pi) = -2$

19. $|z| = 5$, $\theta = \arg(z) = \pi \Rightarrow \sin \theta = 3/5$, $\cos \theta = 4/5$
 $z = 4 + 3i$

20. $|z| = 1$, $\arg(z) = \frac{3\pi}{4} \Rightarrow z = \left(\cos \frac{3\pi}{4} + i \sin \frac{3\pi}{4}\right)$
 $\Rightarrow z = -\frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}}i$

21. $|z| = \pi$, $\arg(z) = \frac{\pi}{6} \Rightarrow z = \pi \left(\cos \frac{\pi}{6} + i \sin \frac{\pi}{6}\right)$
 $\Rightarrow z = \frac{\pi\sqrt{3}}{2} + \frac{\pi}{2}i$

22. $|z| = 0 \Rightarrow z = 0$ for any value of $\arg(z)$

23. $|z| = \frac{1}{2}$, $\arg(z) = -\frac{\pi}{3} \Rightarrow z = \frac{1}{2} \left(\cos \frac{\pi}{3} - i \sin \frac{\pi}{3}\right)$
 $\Rightarrow z = \frac{1}{4} - \frac{\sqrt{3}}{4}i$

24. $\overline{5 + 3i} = 5 - 3i$

25. $\overline{-3 - 5i} = -3 + 5i$

26. $\overline{4i} = -4i$

27. $\overline{2 - i} = 2 + i$

28. $|z| = 2$ represents all points on the circle of radius 2 centred at the origin.

29. $|z| \leq 2$ represents all points in the closed disk of radius 2 centred at the origin.

30. $|z - 2i| \leq 3$ represents all points in the closed disk of radius 3 centred at the point $2i$.

31. $|z - 3 + 4i| \leq 5$ represents all points in the closed disk of radius 5 centred at the point $3 - 4i$.

32. $\arg(z) = \pi/3$ represents all points on the ray from the origin in the first quadrant, making angle 60° with the positive direction of the real axis.

33. $\pi \leq \arg(z) \leq 7\pi/4$ represents the closed wedge-shaped region in the third and fourth quadrants bounded by the ray from the origin to $-\infty$ on the real axis and the ray from the origin making angle -45° with the positive direction of the real axis.

34. $(2 + 5i) + (3 - i) = 5 + 4i$

35. $i - (3 - 2i) + (7 - 3i) = -3 + 7 + i + 2i - 3i = 4$

36. $(4 + i)(4 - i) = 16 - i^2 = 17$

37. $(1 + i)(2 - 3i) = 2 + 2i - 3i - 3i^2 = 5 - i$

38. $(a + bi)(\overline{2a - bi}) = (a + bi)(2a + bi) = 2a^2 - b^2 + 3abi$

39. $(2 + i)^3 = 8 + 12i + 6i^2 + i^3 = 2 + 11i$

40. $\frac{2 - i}{2 + i} = \frac{(2 - i)^2}{4 - i^2} = \frac{3 - 4i}{5}$

41. $\frac{1 + 3i}{2 - i} = \frac{(1 + 3i)(2 + i)}{4 - i^2} = \frac{-1 + 7i}{5}$

42. $\frac{1 + i}{i(2 + 3i)} = \frac{1 + i}{-3 + 2i} = \frac{(1 + i)(-3 - 2i)}{9 + 4} = \frac{-1 - 5i}{13}$

43. $\frac{(1 + 2i)(2 - 3i)}{(2 - i)(3 + 2i)} = \frac{8 + i}{8 + i} = 1$

44. If $z = x + yi$ and $w = u + vi$, where x , y , u , and v are real, then

$$\begin{aligned}\overline{z + w} &= \overline{x + u + (y + v)i} \\ &= x + u - (y + v)i = x - yi + u - vi = \bar{z} + \bar{w}.\end{aligned}$$

45. Using the fact that $|zw| = |z||w|$, we have

$$\overline{\left(\frac{z}{w}\right)} = \overline{\left(\frac{z\bar{w}}{|w|^2}\right)} = \frac{\bar{z}\bar{\bar{w}}}{|w|^2} = \frac{\bar{z}w}{\bar{w}w} = \frac{\bar{z}}{\bar{w}}.$$

46. $z = 3 + i\sqrt{3} = 2\sqrt{3}\left(\cos\frac{\pi}{6} + i\sin\frac{\pi}{6}\right)$
 $w = -1 + i\sqrt{3} = 2\left(\cos\frac{2\pi}{3} + i\sin\frac{2\pi}{3}\right)$

$$\begin{aligned}zw &= 4\sqrt{3}\left(\cos\frac{5\pi}{6} + i\sin\frac{5\pi}{6}\right) \\ \frac{z}{w} &= \sqrt{3}\left(\cos\frac{-\pi}{2} + i\sin\frac{-\pi}{2}\right) = -i\sqrt{3}\end{aligned}$$

47. $z = -1 + i = \sqrt{2}\left(\cos\frac{3\pi}{4} + i\sin\frac{3\pi}{4}\right)$
 $w = 3i = 3\left(\cos\frac{\pi}{2} + i\sin\frac{\pi}{2}\right)$
 $zw = 3\sqrt{2}\left(\cos\frac{5\pi}{4} + i\sin\frac{5\pi}{4}\right) = -3 - 3i$
 $\frac{z}{w} = \frac{\sqrt{2}}{3}\left(\cos\frac{\pi}{4} + i\sin\frac{\pi}{4}\right) = \frac{1}{3} + \frac{1}{3}i$

48. $\cos(3\theta) + i\sin(3\theta) = (\cos\theta + i\sin\theta)^3$
 $= \cos^3\theta + 3i\cos^2\theta\sin\theta - 3\cos\theta\sin^2\theta - i\sin^3\theta$
 Thus

$$\begin{aligned}\cos(3\theta) &= \cos^3\theta - 3\cos\theta\sin^2\theta = 4\cos^3\theta - 3\cos\theta \\ \sin(3\theta) &= 3\cos^2\theta\sin\theta - \sin^3\theta = 3\sin\theta - 4\sin^3\theta.\end{aligned}$$

49. a) $\bar{z} = 2/z$ can be rewritten $|z|^2 = z\bar{z} = 2$, so is satisfied by all numbers z on the circle of radius $\sqrt{2}$ centred at the origin.

b) $\bar{z} = -2/z$ can be rewritten $|z|^2 = z\bar{z} = -2$, which has no solutions since the square of $|z|$ is nonnegative for all complex z .

50. If $z = w = -1$, then $zw = 1$, so $\sqrt{zw} = 1$. But if we use $\sqrt{z} = \sqrt{-1} = i$ and the same value for $\sqrt{w}$, then $\sqrt{z}\sqrt{w} = i^2 = -1 \neq \sqrt{zw}$.

51. The three cube roots of $-1 = \cos\pi + i\sin\pi$ are of the form $\cos\theta + i\sin\theta$ where $\theta = \pi/3$, $\theta = \pi$, and $\theta = 5\pi/3$. Thus they are

$$\frac{1}{2} + i\frac{\sqrt{3}}{2}, \quad -1, \quad \frac{1}{2} - i\frac{\sqrt{3}}{2}.$$

52. The three cube roots of $-8i = 8\left(\cos\frac{3\pi}{2} + i\sin\frac{3\pi}{2}\right)$ are of the form $2(\cos\theta + i\sin\theta)$ where $\theta = \pi/2$, $\theta = 7\pi/6$, and $\theta = 11\pi/6$. Thus they are

$$2i, \quad -\sqrt{3} - i, \quad \sqrt{3} - i.$$

53. The three cube roots of $-1 + i = \sqrt{2}\left(\cos\frac{3\pi}{4} + i\sin\frac{3\pi}{4}\right)$ are of the form $2^{1/6}(\cos\theta + i\sin\theta)$ where $\theta = \pi/4$, $\theta = 11\pi/12$, and $\theta = 19\pi/12$.

54. The four fourth roots of $4 = 4(\cos 0 + i\sin 0)$ are of the form $\sqrt{2}(\cos\theta + i\sin\theta)$ where $\theta = 0$, $\theta = \pi/2$, π , and $\theta = 3\pi/2$. Thus they are $\sqrt{2}$, $i\sqrt{2}$, $-\sqrt{2}$, and $-i\sqrt{2}$.

55. The equation $z^4 + 1 - i\sqrt{3} = 0$ has solutions that are the four fourth roots of $-1 + i\sqrt{3} = 2\left(\cos\frac{2\pi}{3} + i\sin\frac{2\pi}{3}\right)$. Thus they are of the form $2^{1/4}(\cos\theta + i\sin\theta)$, where $\theta = \pi/6$, $2\pi/3$, $7\pi/6$, and $5\pi/3$. They are the complex numbers

$$\pm 2^{1/4}\left(\frac{\sqrt{3}}{2} + \frac{1}{2}i\right), \quad \pm 2^{1/4}\left(\frac{1}{2} - \frac{\sqrt{3}}{2}i\right).$$

56. The equation $z^5 + a^5 = 0$ ($a > 0$) has solutions that are the five fifth roots of $-a^5 = a(\cos \pi + i \sin \pi)$; they are of the form $a(\cos \theta + i \sin \theta)$, where $\theta = \pi/5, 3\pi/5, \pi, 7\pi/5$, and $9\pi/5$.

57. The n th roots of unity are

$$\begin{aligned}\omega_1 &= 1 \\ \omega_2 &= \cos \frac{2\pi}{n} + i \sin \frac{2\pi}{n} \\ \omega_3 &= \cos \frac{4\pi}{n} + i \sin \frac{4\pi}{n} = \omega_2^2 \\ \omega_4 &= \cos \frac{6\pi}{n} + i \sin \frac{6\pi}{n} = \omega_2^3 \\ &\vdots \\ \omega_n &= \cos \frac{2(n-1)\pi}{n} + i \sin \frac{2(n-1)\pi}{n} = \omega_2^{n-1}.\end{aligned}$$

Hence

$$\begin{aligned}\omega_1 + \omega_2 + \omega_3 + \cdots + \omega_n &= 1 + \omega_2 + \omega_2^2 + \cdots + \omega_2^{n-1} \\ &= \frac{1 - \omega_2^n}{1 - \omega_2} = \frac{0}{1 - \omega_2} = 0.\end{aligned}$$

Appendix II. Complex Functions (page A-19)

In Solutions 1–12, $z = x + yi$ and $w = u + vi$, where x , y , u , and v are real.

- The function $w = \bar{z}$ transforms the closed rectangle $0 \leq x \leq 1, 0 \leq y \leq 2$ to the closed rectangle $0 \leq u \leq 1, -2 \leq v \leq 0$.
- The function $w = \bar{z}$ transforms the line $x + y = 1$ to the line $u - v = 1$.
- The function $w = z^2$ transforms the closed annular sector $1 \leq |z| \leq 2, \pi/2 \leq \arg(z) \leq 3\pi/4$ to the closed annular sector $1 \leq |w| \leq 4, \pi \leq \arg(w) \leq 3\pi/2$.
- The function $w = z^3$ transforms the closed quarter-circular disk $0 \leq |z| \leq 2, 0 \leq \arg(z) \leq \pi/2$ to the closed three-quarter disk $0 \leq |w| \leq 8, 0 \leq \arg(w) \leq 3\pi/2$.
- The function $w = 1/z = \bar{z}/|z|^2$ transforms the closed quarter-circular disk $0 \leq |z| \leq 2, 0 \leq \arg(z) \leq \pi/2$ to the closed region lying on or outside the circle $|w| = 1/2$ and in the fourth quadrant, that is, having $-\pi/2 \leq \arg(w) \leq 0$.
- The function $w = -iz$ rotates the z -plane -90° , so transforms the wedge $\pi/4 \leq \arg(z) \leq \pi/3$ to the wedge $-\pi/4 \leq \arg(z) \leq -\pi/6$.
- The function $w = \sqrt{z}$ transforms the ray $\arg(z) = -\pi/3$ (that is, $\text{Arg}(z) = 5\pi/3$) to the ray $\arg(w) = 5\pi/6$.

8. The function $w = z^2 = x^2 - y^2 + 2xyi$ transforms the line $x = 1$ to $u = 1 - y^2, v = 2y$, which is the parabola $v^2 = 4 - 4u$ with vertex at $w = 1$, opening to the left.

9. The function $w = z^2 = x^2 - y^2 + 2xyi$ transforms the line $y = 1$ to $u = x^2 - 1, v = 2x$, which is the parabola $v^2 = 4u + 4$ with vertex at $w = -1$ and opening to the right.

10. The function $w = 1/z = (x - yi)/(x^2 + y^2)$ transforms the line $x = 1$ to the curve given parametrically by

$$u = \frac{1}{1 + y^2}, \quad v = \frac{-y}{1 + y^2}.$$

This curve is, in fact, a circle,

$$u^2 + v^2 = \frac{1 + y^2}{(1 + y^2)^2} = u,$$

with centre $w = 1/2$ and radius $1/2$.

- The function $w = e^z = e^x \cos y + i e^x \sin y$ transforms the horizontal strip $-\infty < x < \infty, \pi/4 \leq y \leq \pi/2$ to the wedge $\pi/4 \leq \arg(w) \leq \pi/2$, or, equivalently, $u \geq 0, v \geq u$.
- The function $w = e^{iz} = e^{-y}(\cos x + i \sin x)$ transforms the vertical half-strip $0 < x < \pi/2, 0 < y < \infty$ to the first-quadrant part of the unit open disk $|w| = e^{-y} < 1, 0 < \arg(w) = x < \pi/2$, that is $u > 0, v > 0, u^2 + v^2 < 1$.
- $f(z) = z^2 = (x + yi)^2 = x^2 - y^2 + 2xyi$
 $u = x^2 - y^2, \quad v = 2xy$
 $\frac{\partial u}{\partial x} = 2x = \frac{\partial v}{\partial y}, \quad \frac{\partial u}{\partial y} = -2y = -\frac{\partial v}{\partial x}$
 $f'(z) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} = 2x + 2yi = 2z.$
- $f(z) = z^3 = (x + yi)^3 = x^3 - 3xy^2 + (3x^2y - y^3)i$
 $u = x^3 - 3xy^2, \quad v = 3x^2y - y^3$
 $\frac{\partial u}{\partial x} = 3(x^2 - y^2) = \frac{\partial v}{\partial y}, \quad \frac{\partial u}{\partial y} = -6xy = -\frac{\partial v}{\partial x}$
 $f'(z) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} = 3(x^2 - y^2 + 2xyi) = 3z^2.$
- $f(z) = \frac{1}{z} = \frac{x - yi}{x^2 + y^2}$
 $u = \frac{x}{x^2 + y^2}, \quad v = \frac{-y}{x^2 + y^2}$
 $\frac{\partial u}{\partial x} = \frac{y^2 - x^2}{(x^2 + y^2)^2} = \frac{\partial v}{\partial y}, \quad \frac{\partial u}{\partial y} = \frac{-2xy}{(x^2 + y^2)^2} = -\frac{\partial v}{\partial x}$
 $f'(z) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} = \frac{-(x^2 - y^2) + 2xyi}{(x^2 + y^2)^2} = \frac{-(\bar{z})^2}{(z\bar{z})^2} = -\frac{1}{z^2}.$

$$\begin{aligned}
 16. \quad f(z) &= e^{z^2} = e^{x^2-y^2}(\cos(2xy) + i \sin(2xy)) \\
 u &= e^{x^2-y^2} \cos(2xy), \quad v = e^{x^2-y^2} \sin(2xy) \\
 \frac{\partial u}{\partial x} &= e^{x^2-y^2} (2x \cos(2xy) - 2y \sin(2xy)) = \frac{\partial v}{\partial y} \\
 \frac{\partial u}{\partial y} &= -e^{x^2-y^2} (2y \cos(2xy) + 2x \sin(2xy)) = -\frac{\partial v}{\partial x} \\
 f'(z) &= \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} \\
 &= e^{x^2-y^2} [2x \cos(2xy) - 2y \sin(2xy) \\
 &\quad + i(2y \cos(2xy) + 2x \sin(2xy))] \\
 &= (2x + 2yi)e^{x^2-y^2}(\cos(2xy) + i \sin(2xy)) = 2ze^{z^2}.
 \end{aligned}$$

$$17. \quad e^{yi} = \cos y + i \sin y \text{ (for real } y\text{). Replacing } y \text{ by } -y, \text{ we get } e^{-yi} = \cos y - i \sin y \text{ (since } \cos \text{ is even and } \sin \text{ is odd). Adding and subtracting these two formulas gives}$$

$$e^{yi} + e^{-yi} = 2 \cos y, \quad e^{yi} - e^{-yi} = 2i \sin y.$$

$$\text{Thus } \cos y = \frac{e^{yi} + e^{-yi}}{2} \text{ and } \sin y = \frac{e^{yi} - e^{-yi}}{2i}.$$

$$\begin{aligned}
 18. \quad e^{z+2\pi i} &= e^x(\cos(y+2\pi) + i \sin(y+2\pi)) \\
 &= e^x(\cos y + i \sin y) = e^z.
 \end{aligned}$$

Thus e^z is periodic with period $2\pi i$. So is $e^{-z} = 1/e^z$. Since $e^{i(z+2\pi)} = e^{zi+2\pi i} = e^{zi}$, therefore e^{zi} and also e^{-zi} are periodic with period 2π . Hence

$$\cos z = \frac{e^{zi} + e^{-zi}}{2} \text{ and } \sin z = \frac{e^{zi} - e^{-zi}}{2i}$$

are periodic with period 2π , and

$$\cosh z = \frac{e^z + e^{-z}}{2} \text{ and } \sinh z = \frac{e^z - e^{-z}}{2}$$

are periodic with period $2\pi i$.

$$\begin{aligned}
 19. \quad \frac{d}{dz} \cos z &= \frac{d}{dz} \frac{e^{zi} + e^{-zi}}{2} = \frac{ie^{zi} - e^{-zi}}{2} = -\sin z \\
 \frac{d}{dz} \sin z &= \frac{d}{dz} \frac{e^{zi} - e^{-zi}}{2i} = \frac{ie^{zi} + e^{-zi}}{2i} = \cos z \\
 \frac{d}{dz} \cosh z &= \frac{d}{dz} \frac{e^z + e^{-z}}{2} = \frac{e^z - e^{-z}}{2} = \sinh z \\
 \frac{d}{dz} \sinh z &= \frac{d}{dz} \frac{e^z - e^{-z}}{2} = \frac{e^z + e^{-z}}{2} = \cosh z
 \end{aligned}$$

$$\begin{aligned}
 20. \quad \cosh(iz) &= \frac{e^{iz} + e^{-iz}}{2} = \cos z \\
 -i \sinh(iz) &= \frac{1}{i} \frac{e^{iz} - e^{-iz}}{2} = \sin z \\
 \cos(iz) &= \frac{e^{-z} + e^z}{2} = \cosh z \\
 \sin(iz) &= \frac{e^{-z} - e^z}{2i} = i \frac{-e^{-z} + e^z}{2} = i \sinh z
 \end{aligned}$$

$$\begin{aligned}
 21. \quad \cos z &= 0 \Leftrightarrow e^{zi} = -e^{-zi} \Leftrightarrow e^{2zi} = -1 \\
 &\Leftrightarrow e^{-2y}[\cos(2x) + i \sin(2x)] = -1 \\
 &\Leftrightarrow \sin(2x) = 0, \quad e^{-2y} \cos(2x) = -1 \\
 &\Leftrightarrow y = 0, \quad \cos(2x) = -1 \\
 &= \Leftrightarrow y = 0, \quad x = \pm \frac{\pi}{2}, \pm \frac{3\pi}{2}, \dots
 \end{aligned}$$

Thus the only complex zeros of $\cos z$ are its real zeros at $z = (2n+1)\pi/2$ for integers n .

$$\begin{aligned}
 22. \quad \sin z &= 0 \Leftrightarrow e^{zi} = e^{-zi} \Leftrightarrow e^{2zi} = 1 \\
 &\Leftrightarrow e^{-2y}[\cos(2x) + i \sin(2x)] = 1 \\
 &\Leftrightarrow \sin(2x) = 0, \quad e^{-2y} \cos(2x) = 1 \\
 &\Leftrightarrow y = 0, \quad \cos(2x) = 1 \\
 &= \Leftrightarrow y = 0, \quad x = 0, \pm\pi, \pm 2\pi, \dots
 \end{aligned}$$

Thus the only complex zeros of $\sin z$ are its real zeros at $z = n\pi$ for integers n .

$$\begin{aligned}
 23. \quad \text{By Exercises 20 and 21, } \cosh z &= 0 \text{ if and only if } \cos(iz) = 0, \text{ that is, if and only if } z = (2n+1)\pi i/2 \\
 &\text{for integer } n. \\
 \text{Similarly, } \sinh z &= 0 \text{ if and only if } \sin(iz) = 0, \text{ that is, if and only if } z = n\pi i \text{ for integer } n.
 \end{aligned}$$

$$\begin{aligned}
 24. \quad e^z &= e^{x+yi} = e^x \cos y + i e^x \sin y \\
 e^{-z} &= e^{-x-yi} = e^{-x} \cos y - i e^{-x} \sin y \\
 \cosh z &= \frac{e^z + e^{-z}}{2} = \frac{e^x + e^{-x}}{2} \cos y + i \frac{e^x - e^{-x}}{2} \sin y \\
 &= \cosh x \cos y + i \sinh x \sin y \\
 \operatorname{Re}(\cosh z) &= \cosh x \cos y, \quad \operatorname{Im}(\cosh z) = \sinh x \sin y.
 \end{aligned}$$

$$\begin{aligned}
 25. \quad \sinh z &= \frac{e^z - e^{-z}}{2} = \frac{e^x - e^{-x}}{2} \cos y + i \frac{e^x + e^{-x}}{2} \sin y \\
 &= \sinh x \cos y + i \cosh x \sin y \\
 \operatorname{Re}(\sinh z) &= \sinh x \cos y, \quad \operatorname{Im}(\sinh z) = \cosh x \sin y.
 \end{aligned}$$

$$\begin{aligned}
 26. \quad e^{iz} &= e^{-y+xi} = e^{-y} \cos x + i e^{-y} \sin x \\
 e^{-iz} &= e^{y-xi} = e^y \cos x - i e^y \sin x \\
 \cos z &= \frac{e^{iz} + e^{-iz}}{2} = \frac{e^{-y} + e^y}{2} \cos x + i \frac{e^{-y} - e^y}{2} \sin x \\
 &= \cos x \cosh y - i \sin x \sinh y \\
 \operatorname{Re}(\cos z) &= \cos x \cosh y, \quad \operatorname{Im}(\cos z) = -\sin x \sinh y \\
 \sin z &= \frac{e^{iz} - e^{-iz}}{2i} = \frac{e^{-y} - e^y}{2i} \cos x + i \frac{e^{-y} + e^y}{2i} \sin x \\
 &= \sin x \cosh y + i \cos x \sinh y \\
 \operatorname{Re}(\sin z) &= \sin x \cosh y, \quad \operatorname{Im}(\sin z) = \cos x \sinh y.
 \end{aligned}$$

$$27. \quad z^2 + 2iz = 0 \Rightarrow z = 0 \text{ or } z = -2i$$

$$\begin{aligned}
 28. \quad z^2 - 2z + i &= 0 \Rightarrow (z-1)^2 = 1-i \\
 &= \sqrt{2} \left(\cos \frac{7\pi}{4} + i \sin \frac{7\pi}{4} \right) \\
 &\Rightarrow z = 1 \pm 2^{1/4} \left(\cos \frac{7\pi}{8} + i \sin \frac{7\pi}{8} \right)
 \end{aligned}$$

$$29. \quad z^2 + 2z + 5 = 0 \Rightarrow (z + 1)^2 = -4 \\ \Rightarrow z = -1 \pm 2i$$

$$30. \quad z^2 - 2iz - 1 = 0 \Rightarrow (z - i)^2 = 0 \\ \Rightarrow z = i \quad (\text{double root})$$

$$31. \quad z^3 - 3iz^2 - 2z = z(z^2 - 3iz - 2) = 0 \\ \Rightarrow z = 0 \text{ or } z^2 - 3iz - 2 = 0 \\ \Rightarrow z = 0 \text{ or } \left(z - \frac{3}{2}i\right)^2 = -\frac{1}{4} \\ \Rightarrow z = 0 \text{ or } z = \left(\frac{3}{2} \pm \frac{1}{2}\right)i \\ \Rightarrow z = 0 \text{ or } z = i \text{ or } z = 2i$$

$$32. \quad z^4 - 2z^2 + 4 = 0 \Rightarrow (z^2 - 1)^2 = -3 \\ z^2 = 1 - i\sqrt{3} \quad \text{or} \quad z^2 = 1 + i\sqrt{3} \\ z^2 = 2 \left(\cos \frac{5\pi}{3} + i \sin \frac{5\pi}{3} \right), \quad z^2 = 2 \left(\cos \frac{\pi}{3} + i \sin \frac{\pi}{3} \right) \\ z = \pm \sqrt{2} \left(\cos \frac{5\pi}{6} + i \sin \frac{5\pi}{6} \right), \quad \text{or} \\ z = \pm \sqrt{2} \left(\cos \frac{\pi}{6} + i \sin \frac{\pi}{6} \right) \\ z = \pm \left(\sqrt{\frac{3}{2}} - \frac{i}{\sqrt{2}} \right), \quad z = \pm \left(\sqrt{\frac{3}{2}} + \frac{i}{\sqrt{2}} \right)$$

$$33. \quad z^4 + 1 = 0 \Rightarrow z^2 = i \text{ or } z^2 = -i \\ \Rightarrow z = \pm \frac{1+i}{\sqrt{2}}, \quad z = \pm \frac{1-i}{\sqrt{2}} \\ z^4 + 1 = \left(z - \frac{1+i}{\sqrt{2}}\right) \left(z - \frac{1-i}{\sqrt{2}}\right) \\ \times \left(z + \frac{1+i}{\sqrt{2}}\right) \left(z + \frac{1-i}{\sqrt{2}}\right) \\ = \left(\left[z - \frac{1}{\sqrt{2}}\right]^2 + \frac{1}{2}\right) \left(\left[z + \frac{1}{\sqrt{2}}\right]^2 + \frac{1}{2}\right) \\ = (z^2 - \sqrt{2}z + 1)(z^2 + \sqrt{2}z + 1)$$

$$34. \quad \text{Since } P(z) = z^4 - 4z^3 + 12z^2 - 16z + 16 \text{ has real} \\ \text{coefficients, if } z_1 = 1 - \sqrt{3}i \text{ is a zero of } P(z), \text{ then so is} \\ \overline{z_1}. \text{ Now}$$

$$(z - z_1)(z - \overline{z_1}) = (z - 1)^2 + 3 = z^2 - 2z + 4.$$

By long division (details omitted) we discover that

$$\frac{z^4 - 4z^3 + 12z^2 - 16z + 16}{z^2 - 2z + 4} = z^2 - 2z + 4.$$

Thus z_1 and $\overline{z_1}$ are both *double zeros* of $P(z)$. These are the only zeros.

$$35. \quad \text{Since } P(z) = z^5 + 3z^4 + 4z^3 + 4z^2 + 3z + 1 \text{ has real} \\ \text{coefficients, if } z_1 = i \text{ is a zero of } P(z), \text{ then so is} \\ z_2 = -i. \text{ Now}$$

$$(z - z_1)(z - z_2) = (z - i)(z + i) = z^2 + 1.$$

By long division (details omitted) we discover that

$$\frac{z^5 + 3z^4 + 4z^3 + 4z^2 + 3z + 1}{z^2 + 1} = z^3 + 3z^2 + 3z + 1 \\ = (z + 1)^3.$$

Thus $P(z)$ has the five zeros: $i, -i, -1, -1, \text{ and } -1$.

$$36. \quad \text{Since } P(z) = z^5 - 2z^4 - 8z^3 + 8z^2 + 31z - 30 \text{ has real} \\ \text{coefficients, if } z_1 = -2 + i \text{ is a zero of } P(z), \text{ then so is} \\ z_2 = -2 - i. \text{ Now}$$

$$(z - z_1)(z - z_2) = z^2 + 4z + 5.$$

By long division (details omitted) we discover that

$$\frac{z^5 - 2z^4 - 8z^3 + 8z^2 + 31z - 30}{z^2 + 4z + 5} \\ = z^3 - 6z^2 + 11z - 6.$$

Observe that $z_3 = 1$ is a zero of $z^3 - 6z^2 + 11z - 6$. By long division again:

$$\frac{z^3 - 6z^2 + 11z - 6}{z - 1} = z^2 - 5z + 6 = (z - 2)(z - 3).$$

Hence $P(z)$ has the five zeros $-2 + i, -2 - i, 1, 2, \text{ and } 3$.

$$37. \quad \text{If } w = z^4 + z^3 - 2iz - 3 \text{ and } |z| = 2, \text{ then } |z^4| = 16 \text{ and}$$

$$|w - z^4| = |z^3 - 2iz - 3| \leq 8 + 4 + 3 = 15 < 16.$$

By the mapping principle described in the proof of Theorem 2, the image in the w -plane of the circle $|z| = 2$ is a closed curve that winds around the origin the same number of times that the image of z^4 does, namely 4 times.

Appendix III. Continuous Functions (page A-25)

1. To be proved: If $a < b < c$, $f(x) \leq g(x)$ for $a \leq x \leq c$, $\lim_{x \rightarrow b} f(x) = L$, and $\lim_{x \rightarrow b} g(x) = M$, then $L \leq M$.

Proof: Suppose, to the contrary, that $L > M$. Let $\epsilon = (L - M)/3$, so $\epsilon > 0$. There exist numbers $\delta_1 > 0$ and $\delta_2 > 0$ such that if $a \leq x \leq b$, then

$$\begin{aligned} |x - b| < \delta_1 &\Rightarrow |f(x) - L| < \epsilon \\ |x - b| < \delta_2 &\Rightarrow |g(x) - M| < \epsilon. \end{aligned}$$

Thus if $|x - b| < \delta = \min\{\delta_1, \delta_2, b - a, c - b\}$, then

$$f(x) - g(x) > L - \epsilon - M - \epsilon = L - M - 2\epsilon = \frac{L - M}{3} > 0.$$

This contradicts the fact that $f(x) \leq g(x)$ on $[a, b]$. Therefore $L \leq M$.

2. To be proved: If $f(x) \leq K$ on $[a, b)$ and $(b, c]$, and if $\lim_{x \rightarrow b} f(x) = L$, then $L \leq K$.

Proof: If $L > K$, then let $\epsilon = (L - K)/2$; thus $\epsilon > 0$. There exists $\delta > 0$ such that $\delta < b - a$ and $\delta < c - b$, and such that if $0 < |x - b| < \delta$, then $|f(x) - L| < \epsilon$. In this case

$$f(x) > L - \epsilon = L - \frac{L - K}{2} > K,$$

which contradicts the fact that $f(x) \leq K$ on $[a, b)$ and $(b, c]$. Therefore $L \leq K$.

3. Let $\epsilon > 0$ be given. Let $\delta = \epsilon^{1/r}$, ($r > 0$). Then

$$0 < x < \delta \Rightarrow 0 < x^r < \delta^r = \epsilon.$$

Thus $\lim_{x \rightarrow 0^+} x^r = 0$.

4. a) Let $f(x) = C$, $g(x) = x$. Let $\epsilon > 0$ be given and let $\delta = \epsilon$. For any real number x , if $|x - a| < \delta$, then

$$\begin{aligned} |f(x) - f(a)| &= |C - C| = 0 < \epsilon, \\ |g(x) - g(a)| &= |x - a| < \delta = \epsilon. \end{aligned}$$

Thus $\lim_{x \rightarrow a} f(x) = f(a)$ and $\lim_{x \rightarrow a} g(x) = g(a)$, and f and g are both continuous at every real number a .

5. A polynomial is constructed by adding and multiplying finite numbers of functions of the type of f and g in Exercise 4. By Theorem 1(a), such sums and products are continuous everywhere, since their components have been shown to be continuous everywhere.
6. If P and Q are polynomials, they are continuous everywhere by Exercise 5. If $Q(a) \neq 0$, then $\lim_{x \rightarrow a} \frac{P(x)}{Q(x)} = \frac{P(a)}{Q(a)}$ by Theorem 1(a). Hence P/Q is continuous everywhere except at the zeros of Q .

7. Suppose n is a positive integer and $a > 0$. Let $\epsilon > 0$ be given. Let $b = a^{1/n}$, and let $\delta = \min\{a(1 - 2^{-n}), b^{n-1}\epsilon\}$. If $|x - a| < \delta$, then $x > a/2^n$, and if $y = x^{1/n}$, then $y > b/2$. Thus

$$\begin{aligned} |x^{1/n} - a^{1/n}| &= |y - b| \\ &= \frac{|y^n - b^n|}{y^{n-1} + y^{n-2}b + \cdots + b^{n-1}} \\ &< \frac{|x - a|}{b^{n-1}} < \frac{b^{n-1}\epsilon}{b^{n-1}} = \epsilon. \end{aligned}$$

Thus $\lim_{x \rightarrow a} x^{1/n} = a^{1/n}$, and $x^{1/n}$ is continuous at $x = a$.

8. By Exercise 5, x^m is continuous everywhere. By Exercise 7, $x^{1/n}$ is continuous at each $a > 0$. Thus for $a > 0$ we have

$$\begin{aligned} \lim_{x \rightarrow a} x^{m/n} &= \lim_{x \rightarrow a} (x^{1/n})^m = \left(\lim_{x \rightarrow a} x^{1/n} \right)^m \\ &= (a^{1/n})^m = a^{m/n}, \end{aligned}$$

and $x^{m/n}$ is continuous at each positive number.

9. If m and n are integers and n is odd, then $(-x)^{m/n} = cx^{m/n}$, where $c = (-1)^{m/n}$ is either -1 or 1 depending on the parity of m . Since $x^{m/n}$ is continuous at each positive number a , so is $cx^{m/n}$. Thus $(-x)^{m/n}$ is continuous at each positive number, and $x^{m/n}$ is continuous at each negative number.

If $r = m/n > 0$, then $\lim_{x \rightarrow 0^+} x^r = 0$ by Exercise 3. Hence $\lim_{x \rightarrow 0^-} x^r = (-1)^r \lim_{x \rightarrow 0^+} x^r = 0$, also. Therefore $\lim_{x \rightarrow 0} x^r = 0$, and x^r is continuous at $x = 0$.

10. Let $\epsilon > 0$ be given. Let $\delta = \epsilon$. If a is any real number then

$$||x| - |a|| \leq |x - a| < \epsilon \quad \text{if} \quad |x - a| < \delta.$$

Thus $\lim_{x \rightarrow a} |x| = |a|$, and the absolute value function is continuous at every real number.

11. By the definition of $\sin$, $P_t = (\cos t, \sin t)$, and $P_a = (\cos a, \sin a)$ are two points on the unit circle $x^2 + y^2 = 1$. Therefore

$$\begin{aligned} |t - a| &= \text{length of the arc from } P_t \text{ to } P_a \\ &> \text{length of the chord from } P_t \text{ to } P_a \\ &= \sqrt{(\cos t - \cos a)^2 + (\sin t - \sin a)^2}. \end{aligned}$$

If $\epsilon > 0$ is given, and $|t - a| < \delta = \epsilon$, then the above inequality implies that

$$\begin{aligned} |\cos t - \cos a| &\leq |t - a| < \epsilon, \\ |\sin t - \sin a| &\leq |t - a| < \epsilon. \end{aligned}$$

Thus $\sin$ is continuous everywhere.

12. The proof that $\cos$ is continuous everywhere is almost identical to that for $\sin$ in Exercise 11.

13. Let $a > 0$ and $\epsilon > 0$. Let $\delta = \min \left\{ \frac{a}{2}, \frac{\epsilon a}{2} \right\}$.

If $|x - a| < \delta$, then $x > \frac{a}{2}$, so $\frac{1}{t} < \frac{2}{a}$ whenever t is between a and x . Thus

$$\begin{aligned} |\ln x - \ln a| &= \text{area under } y = \frac{1}{t} \text{ between } t = a \text{ and } t = x \\ &< \frac{2}{a}|x - a| < \frac{2}{a} \frac{\epsilon a}{2} = \epsilon. \end{aligned}$$

Thus $\lim_{x \rightarrow a} \ln x = \ln a$, and $\ln$ is continuous at each point a in its domain $(0, \infty)$.

14. Let a be any real number, and let $\epsilon > 0$ be given. Assume (making ϵ smaller if necessary) that $\epsilon < e^a$. Since

$$\ln \left(1 - \frac{\epsilon}{e^a} \right) + \ln \left(1 + \frac{\epsilon}{e^a} \right) = \ln \left(1 - \frac{\epsilon^2}{e^{2a}} \right) < 0,$$

we have $\ln \left(1 + \frac{\epsilon}{e^a} \right) < -\ln \left(1 - \frac{\epsilon}{e^a} \right)$.

Let $\delta = \ln \left(1 + \frac{\epsilon}{e^a} \right)$. If $|x - a| < \delta$, then

$$\begin{aligned} \ln \left(1 - \frac{\epsilon}{e^a} \right) &< x - a < \ln \left(1 + \frac{\epsilon}{e^a} \right) \\ 1 - \frac{\epsilon}{e^a} &< e^{x-a} < 1 + \frac{\epsilon}{e^a} \\ |e^{x-a} - 1| &< \frac{\epsilon}{e^a} \\ |e^x - e^a| &= e^a |e^{x-a} - 1| < \epsilon. \end{aligned}$$

Thus $\lim_{x \rightarrow a} e^x = e^a$ and e^x is continuous at every point a in its domain.

15. Suppose $a \leq x_n \leq b$ for each n , and $\lim x_n = L$. Then $a \leq L \leq b$ by Theorem 3. Let $\epsilon > 0$ be given. Since f is continuous on $[a, b]$, there exists $\delta > 0$ such that if $a \leq x \leq b$ and $|x - L| < \delta$ then $|f(x) - f(L)| < \epsilon$. Since $\lim x_n = L$, there exists an integer N such that if $n \geq N$ then $|x_n - L| < \delta$. Hence $|f(x_n) - f(L)| < \epsilon$ for such n . Therefore $\lim(f(x_n)) = f(L)$.

16. Let $g(t) = \frac{t}{1 + |t|}$. For $t \neq 0$ we have

$$g'(t) = \frac{1 + |t| - t \operatorname{sgn} t}{(1 + |t|)^2} = \frac{1 + |t| - |t|}{(1 + |t|)^2} = \frac{1}{(1 + |t|)^2} > 0.$$

If $t = 0$, g is also differentiable, and has derivative 1:

$$g'(0) = \lim_{h \rightarrow 0} \frac{g(h) - g(0)}{h} = \lim_{h \rightarrow 0} \frac{1}{1 + |h|} = 1.$$

Thus g is continuous and increasing on $\mathbb{R}$.

If f is continuous on $[a, b]$, then

$$h(x) = g(f(x)) = \frac{f(x)}{1 + |f(x)|}$$

is also continuous there, being the composition of continuous functions. Also, $h(x)$ is bounded on $[a, b]$, since

$$|g(f(x))| \leq \frac{|f(x)|}{1 + |f(x)|} \leq 1.$$

By assumption in this problem, $h(x)$ must assume maximum and minimum values; there exist c and d in $[a, b]$ such that

$$g(f(c)) \leq g(f(x)) \leq g(f(d))$$

for all x in $[a, b]$. Since g is increasing, so is its inverse g^{-1} . Therefore

$$f(c) \leq f(x) \leq f(d)$$

for all x in $[a, b]$, and f is bounded on that interval.

Appendix IV. The Riemann Integral (page A-30)

1. $f(x) = \begin{cases} 1 & \text{if } 0 \leq x \leq 1 \\ 0 & \text{if } 1 < x \leq 2 \end{cases}$

Let $0 < \epsilon < 1$. Let $P = \{0, 1 - \frac{\epsilon}{3}, 1 + \frac{\epsilon}{3}, 2\}$. Then

$$L(f, P) = 1 \left(1 - \frac{\epsilon}{3} \right) + 0 + 0 = 1 - \frac{\epsilon}{3}$$

$$U(f, P) = 1 \left(1 - \frac{\epsilon}{3} \right) + 1 \left(\frac{2\epsilon}{3} \right) + 0 = 1 + \frac{\epsilon}{3}.$$

Since $U(f, P) - L(f, P) < \epsilon$, f is integrable on $[0, 2]$.

Since $L(f, P) < 1 < U(f, P)$ for every ϵ , therefore

$$\int_0^2 f(x) dx = 1.$$

2. $f(x) = \begin{cases} 1 & \text{if } x = 1/n \quad (n = 1, 2, 3, \dots) \\ 0 & \text{otherwise} \end{cases}$

If P is any partition of $[0, 1]$ then $L(f, P) = 0$. Let

$0 < \epsilon \leq 2$. Let N be an integer such that

$N + 1 > \frac{2}{\epsilon} \geq N$. A partition P of $[0, 1]$

can be constructed so that the first two points of P

are 0 and $\frac{\epsilon}{2}$, and such that each of the N points $\frac{1}{n}$

($n = 1, 2, 3, \dots, n$) lies in a subinterval of P having

length at most $\frac{\epsilon}{2N}$. Since every number $\frac{1}{n}$ with n a positive integer lies either in $\left[0, \frac{\epsilon}{2}\right]$ or one of these other

N subintervals of P , and since $\max f(x) = 1$ for these subintervals and $\max f(x) = 0$ for all other subintervals

of P , therefore $U(f, P) \leq \frac{\epsilon}{2} + N \frac{\epsilon}{2N} = \epsilon$. By Theorem

3, f is integrable on $[0, 1]$. Evidently

$$\int_0^1 f(x) dx = \text{least upper bound } L(f, P) = 0.$$

3. $f(x) = \begin{cases} 1/n & \text{if } x = m/n \text{ in lowest terms} \\ 0 & \text{otherwise} \end{cases}$

Clearly $L(f, P) = 0$ for every partition P of $[0, 1]$. Let $\epsilon > 0$ be given. To show that f is integrable we must exhibit a partition P for which $U(f, P) < \epsilon$. We can assume $\epsilon < 1$. Choose a positive integer N such that $2/N < \epsilon$. There are only finitely many integers n such that $1 \leq n \leq N$. For each such n , there are only finitely many integers m such that $0 \leq m/n \leq 1$. Therefore there are only finitely many points x in $[0, 1]$ where $f(x) > \epsilon/2$. Let P be a partition of $[0, 1]$ such that all these points are contained in subintervals of the partition having total length less than $\epsilon/2$. Since $f(x) \leq 1$ on these subintervals, and $f(x) < \epsilon/2$ on all other subintervals P , therefore $U(f, P) \leq 1 \times (\epsilon/2) + (\epsilon/2) \times 1 = \epsilon$, and f is integrable on $[0, 1]$. Evidently $\int_0^1 f(x) dx = 0$, since all lower sums are 0.

4. Suppose, to the contrary, that $I_* > I^*$. Let $\epsilon = \frac{I_* - I^*}{3}$, so $\epsilon > 0$. By the definition of I_* and I^* , there exist partitions P_1 and P_2 of $[a, b]$, such that $L(f, P_1) \geq I_* - \epsilon$ and $U(f, P_2) \leq I^* + \epsilon$. By Theorem 2, $L(f, P_1) \leq U(f, P_2)$, so

$$3\epsilon = I_* - I^* \leq L(f, P_1) + \epsilon - U(f, P_2) + \epsilon \leq 2\epsilon.$$

Since $\epsilon > 0$, it follows that $3 \leq 2$. This contradiction shows that we must have $I_* \leq I^*$.

5. Theorem 3 of Section 6.4: Proofs of parts (c)–(h).

- c) Multiplying a function by a constant multiplies all its Riemann sums by the same constant. If the constant is positive, upper and lower sums remain upper and lower; if the constant is negative upper sums become lower and vice versa. Therefore

$$\int_a^b Af(x) dx = A \int_a^b f(x) dx.$$

It therefore remains to be proved only that the integral of a sum of functions is the sum of the integrals. Suppose that

$$\int_a^b f(x) dx = I, \quad \text{and} \quad \int_a^b g(x) dx = J.$$

If $\epsilon > 0$, then there exist partitions P_1 and P_2 of $[a, b]$ such that

$$\begin{aligned} U(f, P_1) - \frac{\epsilon}{2} &\leq I < L(f, P_1) + \frac{\epsilon}{2} \\ U(g, P_2) - \frac{\epsilon}{2} &\leq J < L(g, P_2) + \frac{\epsilon}{2}. \end{aligned}$$

Let P be the common refinement of P_1 and P_2 . Then the above inequalities hold with P replacing P_1 and P_2 . If $m_1 \leq f(x) \leq M_1$ and $m_2 \leq g(x) \leq M_2$ on any interval, then $m_1 + m_2 \leq f(x) + g(x) \leq M_1 + M_2$ there. It follows that

$$\begin{aligned} U(f + g, P) &\leq U(f, P) + U(g, P), \\ L(f, P) + L(g, P) &\leq L(f + g, P). \end{aligned}$$

Therefore

$$U(f + g, P) - \epsilon \leq I + J \leq L(f + g, P) + \epsilon.$$

Hence $\int_a^b (f(x) + g(x)) dx = I + J$.

- d) Assume $a < b < c$; the other cases are similar. Let $\epsilon > 0$. If

$$\int_a^b f(x) dx = I, \quad \text{and} \quad \int_b^c f(x) dx = J,$$

then there exist partitions P_1 of $[a, b]$, and P_2 of $[b, c]$ such that

$$\begin{aligned} L(f, P_1) &\leq I < L(f, P_1) + \frac{\epsilon}{2} \\ L(f, P_2) &\leq J < L(f, P_2) + \frac{\epsilon}{2} \end{aligned}$$

(with similar inequalities for upper sums). Let P be the partition of $[a, c]$ formed by combining all the subdivision points of P_1 and P_2 . Then

$$L(f, P) = L(f, P_1) + L(f, P_2) \leq I + J < L(f, P) + \epsilon.$$

Similarly, $U(f, P) - \epsilon < I + J \leq U(f, P)$. Therefore

$$\int_a^c f(x) dx = I + J.$$

- e) Let

$$\int_a^b f(x) dx = I, \quad \text{and} \quad \int_a^b g(x) dx = J,$$

where $f(x) \leq g(x)$ on $[a, b]$. We want to show that $I \leq J$. Suppose, to the contrary, that $I > J$. Then there would exist a partition P of $[a, b]$ for which

$$I < L(f, P) + \frac{I - J}{2}, \quad \text{and} \quad U(g, P) - \frac{I - J}{2} < J.$$

Thus $L(f, P) > \frac{I + J}{2} > U(g, P) \geq L(g, P)$.

However, $f(x) \leq g(x)$ on $[a, b]$ implies that $L(f, P) \leq L(g, P)$ for any partition. Thus we have a contradiction, and so $I \leq J$.

- f) Since $-|f(x)| \leq f(x) \leq |f(x)|$ for any x , we have by part (e), if $a \leq b$,

$$-\int_a^b |f(x)| dx \leq \int_a^b f(x) dx \leq \int_a^b |f(x)| dx.$$

$$\text{Therefore } \left| \int_a^b f(x) dx \right| \leq \int_a^b |f(x)| dx.$$

- g) By parts (b), (c) and (d),

$$\begin{aligned} \int_{-a}^a f(x) dx &= \int_{-a}^0 f(x) dx + \int_0^a f(x) dx \\ &= \int_0^a f(-x) dx + \int_0^a f(x) dx \\ &= \int_0^a [f(-x) + f(x)] dx. \end{aligned}$$

If f is odd, the last integral is 0. If f is even, the last integral is $\int_0^a 2f(x) dx$. Thus both (g) and (h) are proved.

6. Let $\epsilon > 0$ be given. Let $\delta = \epsilon^2/2$. Let $0 \leq x \leq 1$ and $0 \leq y \leq 1$. If $x < \epsilon^2/4$ and $y < \epsilon^2/4$ then $|\sqrt{x} - \sqrt{y}| \leq \sqrt{x} + \sqrt{y} < \epsilon$. If $|x - y| < \delta$ and either $x \geq \epsilon^2/4$ or $y \geq \epsilon^2/4$ then

$$|\sqrt{x} - \sqrt{y}| = \frac{|x - y|}{\sqrt{x} + \sqrt{y}} < \frac{2}{\epsilon} \times \frac{\epsilon^2}{2} = \epsilon.$$

Thus $f(x) = \sqrt{x}$ is uniformly continuous on $[0, 1]$.

7. Suppose f is uniformly continuous on $[a, b]$. Taking $\epsilon = 1$ in the definition of uniform continuity, we can find a positive number δ such that $|f(x) - f(y)| < 1$ whenever x and y are in $[a, b]$ and $|x - y| < \delta$. Let N be a positive integer such that $h = (b - a)/N$ satisfies $h < \delta$. If $x_k = a + kh$, ($0 \leq k \leq N$), then each of the subintervals of the partition $P = \{x_0, x_1, \dots, x_N\}$ has length less than δ . Thus

$$|f(x_k) - f(x_{k-1})| < 1 \quad \text{for } 1 \leq k \leq N.$$

By repeated applications of the triangle inequality,

$$|f(x_{k-1}) - f(a)| = |f(x_{k-1}) - f(x_0)| < k - 1.$$

If x is any point in $[a, b]$, then x belongs to one of the intervals $[x_{k-1}, x_k]$, so, by the triangle inequality again,

$$|f(x) - f(a)| \leq |f(x) - f(x_{k-1})| + |f(x_{k-1}) - f(a)| < k \leq N.$$

Thus $|f(x)| < |f(a)| + N$, and f is bounded on $[a, b]$.

8. Suppose that $|f(x)| \leq K$ on $[a, b]$ (where $K > 0$), and that f is integrable on $[a, b]$. Let $\epsilon > 0$ be given, and let $\delta = \epsilon/K$. If x and y belong to $[a, b]$ and $|x - y| < \delta$, then

$$\begin{aligned} |F(x) - F(y)| &= \left| \int_a^x f(t) dt - \int_a^y f(t) dt \right| \\ &= \left| \int_y^x f(t) dt \right| \leq K|x - y| < K\delta = \epsilon. \end{aligned}$$

(See Theorem 3(f) of Section 6.4.) Thus F is uniformly continuous on $[a, b]$.